

# Using Medications Wisely (National Patient Safety Goals 2012)

1. Plural Noun
2. Plural Noun
3. Past Tense Verb
4. Adjective
5. Noun
6. Noun
7. Adverb
8. Plural Noun
9. Noun
10. Adjective
11. Noun
12. Adverb
13. Adverb
14. Adjective
15. Past Tense Verb
16. Adjective
17. Adjective
18. Adjective
19. Adjective
20. Adjective
21. Adjective
22. Plural Noun

23. Verb Ending In Ing

24. Preposition

25. Noun

26. Noun

27. Noun

28. Adjective

29. Adjective

30. Noun

31. Past Tense Verb

32. Verb

33. Noun

34. Adjective

35. Noun

36. Noun

37. Noun

38. Noun

39. Noun

40. Present Tense Verb

# Using Medications Wisely (National Patient Safety Goals 2012)

or other solutions in unlabeled \_\_\_\_\_ or syringes are unidentifiable. Tragic errors, including deaths, have occurred from the removal of medications or solutions from their original containers and placed into \_\_\_\_\_ containers. This practice is commonplace in many organizations and yet it negates the basic principle of \_\_\_\_\_ management.

Unless a \_\_\_\_\_ is prepared and \_\_\_\_\_ administered to a patient by the same individual, it must be appropriately labeled as to its \_\_\_\_\_ strength, diluent, volume and expiration \_\_\_\_\_ and time. For short procedures, such as epidural procedures in the pain management clinic, dating and timing of containers may not be necessary as defined by the hospital. Any medication being prepared by one individual and administered by a \_\_\_\_\_ must be appropriately labeled as described above and verified both \_\_\_\_\_ and \_\_\_\_\_.

\_\_\_\_\_ therapy is a useful therapeutic treatment for a variety of conditions such as atrial fibrillation, heart valve implantation, and venous/arterial thrombotic events. Anticoagulants are also \_\_\_\_\_ with significant patient harm due to \_\_\_\_\_ dosing, \_\_\_\_\_ monitoring and patient \_\_\_\_\_.

Bringing awareness to the appropriate use of anticoagulants will improve patient safety and achieve positive patient outcomes.

The availability of oral \_\_\_\_\_ Adjective \_\_\_\_\_ products, \_\_\_\_\_ Adjective \_\_\_\_\_ syringes, and \_\_\_\_\_ Adjective \_\_\_\_\_ intravenous solutions should be utilized wherever possible to reduce the likelihood of dosing errors. Following approved \_\_\_\_\_ Plural noun \_\_\_\_\_ for initiating and \_\_\_\_\_ Verb ending in ing \_\_\_\_\_ anticoagulant therapy will enhance positive outcomes a?? keeping patient variable response to anticoagulant therapy in mind. Educating the patient regarding his/her therapy is \_\_\_\_\_ Preposition \_\_\_\_\_ to compliance with the therapy and improving their outcome. \_\_\_\_\_ Noun \_\_\_\_\_ should cover \_\_\_\_\_ Noun \_\_\_\_\_ and \_\_\_\_\_ Noun \_\_\_\_\_ interactions with their therapy, laboratory monitoring of their progress, and compliance with the therapy.

Medication reconciliation between \_\_\_\_\_ Adjective \_\_\_\_\_ therapy and \_\_\_\_\_ Adjective \_\_\_\_\_ therapy is gaining more recognition as an important part of the admission process. Any \_\_\_\_\_ Noun \_\_\_\_\_ between what they were taking at home and what they are currently being prescribed can be \_\_\_\_\_ Past tense verb \_\_\_\_\_ prior to medication administration and thereby \_\_\_\_\_ Verb \_\_\_\_\_ patient harm.

Upon \_\_\_\_\_ the patient should receive written information regarding their \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_. This should include \_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
route, \_\_\_\_\_  
\_\_\_\_\_  
and reason for  
\_\_\_\_\_.  
Patients should be encouraged to provide a list of their current medication therapy to their  
primary care physician. \_\_\_\_\_  
\_\_\_\_\_  
in dose, medications, discontinuation of therapy, etc should be  
communicated to the patient and their primary care physician as well.