

Using Medications Wisely (National Patient Safety Goals 2012)

1. Plural Noun
2. Plural Noun
3. Past Tense Verb
4. Adjective
5. Noun
6. Noun
7. Adverb
8. Plural Noun
9. Noun
10. Adjective
11. Noun
12. Adverb
13. Adverb
14. Adjective
15. Past Tense Verb
16. Adjective
17. Adjective
18. Adjective
19. Adjective
20. Adjective
21. Adjective
22. Plural Noun

23. Verb Ending In Ing

24. Preposition

25. Noun

26. Noun

27. Noun

28. Adjective

29. Adjective

30. Noun

31. Past Tense Verb

32. Verb

33. Noun

34. Adjective

35. Noun

36. Noun

37. Noun

38. Noun

39. Noun

40. Present Tense Verb

Using Medications Wisely (National Patient Safety Goals 2012)

or other solutions in unlabeled _____
Plural noun or syringes are unidentifiable. Tragic errors, including deaths, have occurred from the removal of medications or solutions from their original containers and placed into _____
Past tense verb containers. This practice is commonplace in many organizations and yet it negates the basic principle of _____
Adjective _____
Noun management.

Unless a _____
Noun is prepared and _____
Adverb administered to a patient by the same individual, it must be appropriately labeled as to its _____
Plural noun strength, diluent, volume and expiration _____
Noun and time. For short procedures, such as epidural procedures in the pain management clinic, dating and timing of containers may not be necessary as defined by the hospital. Any medication being prepared by one individual and administered by a _____
Adjective _____
Noun must be appropriately labeled as described above and verified both _____
Adverb and _____
Adverb.

Adjective therapy is a useful therapeutic treatment for a variety of conditions such as atrial fibrillation, heart valve implantation, and venous/arterial thrombotic events. Anticoagulants are also _____
Past tense verb with significant patient harm due to _____
Adjective dosing, _____
Adjective monitoring and patient _____
Adjective.

Bringing awareness to the appropriate use of anticoagulants will improve patient safety and achieve positive patient outcomes.

The availability of oral _____
Adjective products, _____
Adjective syringes, and _____
Adjective intravenous solutions should be utilized wherever possible to reduce the likelihood of dosing errors. Following approved _____
Plural noun for initiating and _____
Verb ending in ing anticoagulant therapy will enhance positive outcomes
â?? keeping patient variable response to anticoagulant therapy in mind. Educating the patient regarding his/her therapy is _____
Preposition to compliance with the therapy and improving their outcome. _____
Noun should cover _____
Noun and _____
Noun interactions with their therapy, laboratory monitoring of their progress, and compliance with the therapy.

Medication reconciliation between _____
Adjective therapy and _____
Adjective therapy is gaining more recognition as an important part of the admission process. Any _____
Noun between what they were taking at home and what they are currently being prescribed can be _____
Past tense verb prior to medication administration and thereby _____
Verb patient harm.

Upon _____ the patient should receive written information regarding their _____.
_____. This should include _____ route, _____ and reason for
_____. Patients should be encouraged to provide a list of their current medication therapy to their
primary care physician. _____ in dose, medications, discontinuation of therapy, etc should be
communicated to the patient and their primary care physician as well.